



Investigating the phenology of seaward migration of juvenile brown trout (*Salmo trutta*) in two European populations

Winter, E R.; Tummers, J. S.; Aarestrup, Kim; Baktoft, Henrik; Lucas, M. C.

Published in:
Journal of Applied Phycology

Link to article, DOI:
[10.1007/s10750-016-2720-z](https://doi.org/10.1007/s10750-016-2720-z)

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Winter, E. R., Tummers, J. S., Aarestrup, K., Baktoft, H., & Lucas, M. C. (2016). Investigating the phenology of seaward migration of juvenile brown trout (*Salmo trutta*) in two European populations. *Journal of Applied Phycology*, 775(1), 139-151. <https://doi.org/10.1007/s10750-016-2720-z>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Durham Research Online

Deposited in DRO:

25 July 2017

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Winter, E.R. and Tummers, J.S. and Aarestrup, K. and Baktoft, H. and Lucas, M.C. (2016) 'Investigating the phenology of seaward migration of juvenile brown trout (*Salmo trutta*) in two European populations.', *Hydrobiologia.*, 775 (1). pp. 139-151.

Further information on publisher's website:

<https://doi.org/10.1007/s10750-016-2720-z>

Publisher's copyright statement:

The final publication is available at Springer via <https://doi.org/10.1007/s10750-016-2720-z>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Investigating the phenology of seaward migration of juvenile
brown trout (*Salmo trutta*) in two European populations

Final accepted version, *Hydrobiologia* Feb 2016

E. R. Winter^a, J. S. Tummers^a, K. Aarestrup^b, H. Baktoft^b & M. C. Lucas^a

^a School of Biological and Biomedical Sciences, Durham University, South Road,
Durham, UK, DH1 3LE

^b Technical University of Denmark, Vejlsøvej 39, 8600 Silkeborg

Corresponding author: E. R. Winter: e.r.winter@durham.ac.uk,
ewinter64@yahoo.co.uk

Abstract

Recent evidence supports the existence of a downstream autumn migratory phenotype in juvenile anadromous brown trout (*Salmo trutta*), however the precise timing, extent and ecological significance of such behaviour remains ambiguous. We investigated the phenology of downstream migration of wild juvenile trout using passive integrated transponder (PIT) telemetry over an eight month period in two European rivers; the River Deerness, north-east England, and the River Villestrup, Denmark. The incidence of autumn-winter seaward migration was greater in the Deerness than the Villestrup, with at least 46 % of migrating juveniles detected prior to spring smoltification in the Deerness. Timing of migration was strongly regulated by factors associated with river discharge in both systems. While autumn and spring downstream migrants did not differ in size at the time of tagging in either system, evidence that spring migrants were of better condition, travelled faster (autumn: 11.0 km day⁻¹; spring: 24.3 km day⁻¹) and were more likely to leave the Deerness suggests autumn and spring migrant conspecifics respond to different behavioural motivations. Further investigation into the sex of autumn migrant juveniles, as well as the temporal and geographical variability in the incidence and fitness consequences of autumn emigration by juvenile trout would be beneficial.

Keywords: Autumn migration, smolt, life history, phenotype, phenology, telemetry

Introduction

Individuals of migratory species often exhibit wide variation in their spatial behaviour, varying from local residency to large-scale migration (Nathan et al. 2008; Chapman et al. 2011), including for fishes (Lucas & Baras, 2001). The literature concerning such variation within the Salmonidae is voluminous (reviewed in Dodson et al. 2013). The brown trout, *Salmo trutta* L., exemplifies such phenotypic plasticity with its spatial and temporal continuum of alternative migratory tactics, from locally-resident to potamodromous and anadromous forms (Jonsson & Jonsson, 2011; Boel et al. 2014). Partial and differential migration, at the intrapopulation level, have been illustrated in numerous *S. trutta* populations, notably around Vangsvatnet Lake, Norway (Jonsson, 1985), in the Oir River, France (Cucherousset et al. 2005) and with regard to the extent of marine migration (del Villar-Guerra et al. 2013; Aarestrup et al. 2014).

Despite thorough accounts of behaviour, elements of juvenile salmonid space use and phenology of downstream migration remain poorly explained. It is traditionally considered that juvenile populations of anadromous brown trout and closely related Atlantic salmon, *S. salar*, remain in their native stream from one to several years before descending the river course as smolts in spring (Klemetsen et al. 2003). Therefore, when considering anadromous salmonid population dynamics, the freshwater output of a river typically refers to the production of spring migrants (Ibbotson et al. 2013). Estimates of juvenile salmonid population density are usually carried out in late spring when site fidelity is regarded as strong (Cunjak 1992) and one is typically unable to distinguish mortality from emigration when examining population losses over autumn and winter. Few published tracking or trapping studies have targeted juvenile trout outside spring (but see Jonsson & Jonsson, 2002, 2009; Marine Institute, 2014; Holmes et al. 2014; Taal et al. 2014). For example, in widely cited studies on trout migration, Elliott (1986, 1994) did not investigate evidence that small-scale emigration of Age 1+ trout parr from Black Brows Beck, NW England, appeared to occur year round and represented a high proportion, relative to spring smolt numbers.

Downstream movements of *S. salar* juveniles are well known in autumn (Youngson et al. 1983; Cunjak et al. 1989; Pinder et al. 2007; McGinnity et al. 2007; Jensen et al. 2012; Jonsson & Jonsson, 2014; Taal et al. 2014), although the proportions of populations undertaking those movements, and their significance, are still not fully understood. By contrast, autumn descents of immature brown trout have been less widely recorded, but information is available in Irish (Marine Institute, 2014), Norwegian (Jonsson & Jonsson, 2002, 2009), Baltic (Taal et al. 2014) and New Zealand (Holmes et al. 2014) populations. Based upon population-specific interpretations of phenotype, non-moribund, juvenile salmonid autumn migrants are regarded as pre-smolts, destined to enter the sea (McGinnity et al. 2007), or as comprising potamodromous population components that may be common in larger river systems (Cucherousset et al. 2005). Survival may also differ by population, for example the weakly brackish environment encountered by autumn migrants in the Baltic Sea does not require marine-level osmoregulatory competency for survival (Taal et al. 2014). Alternatively, ice, low river discharge and the absence of an estuary in which to reside are thought to cause poor survival of *S. salar* autumn migrants leaving the River Halselva, Norway and a lack of sympatric *S. trutta* conspecifics (Jensen et al. 2012). Autumn migrant *S. salar* do not appear suitably adapted to seawater (Riley et al. 2008), and sea survival rates of first-time, autumn-emigrant *S. trutta* in Norway are significantly lower than those in spring (Jonsson & Jonsson, 2009), however the viability of these phenotypes has been confirmed with returning *Salmo* adults (Jonsson & Jonsson, 2009; Riley et al. 2009).

Despite enhanced awareness of an autumn downstream migratory phenotype in juvenile salmonids over recent years, our understanding of its ecological significance remains limited and somewhat contested. Recorded incidence of autumn migration has been associated with proximity to the marine environment (Ibbotson et al. 2013), elevated stream discharge (Youngson et al. 1983) and poor overwintering habitat (Riddell & Leggett, 1981), as well as the reproductive motivations of precociously mature male parr (Buck & Youngson, 1982; McGinnity et al. 2007; Jensen et al. 2012). Holmes et al. (2014) suggested that early emigration of larger trout parr from the Rainy River, New

Zealand reflected limited overwintering habitat availability and/or constrained feeding opportunities for individuals with high growth rates. Determining the value of alternative wintering habitats, in terms of trout survival, smolt output, performance in the sea and ultimately their fecundity, is important for river and fisheries managers in directing conservation, regulation and habitat management (Ibbotson et al. 2013), such as when considering the importance of year-round downstream fish passage.

In this study we examined and compared the phenology of downstream migration in two juvenile trout populations in north-east England and Denmark, using Passive Integrated Transponder (PIT) telemetry. We investigated the influence of environmental, subject-specific and tag-site specific variables on the probability of passage past fixed monitoring sites over an eight month period.

Materials and Methods

Study areas

The River Deerness (mean annual discharge $\sim 0.5 \text{ m}^3 \text{ s}^{-1}$ in its lower reaches) is a small tributary (width in study reaches, 2-10 m) of the River Browney in NE England ($54^\circ 44' \text{ N}$, $1^\circ 48' \text{ W}$) and flows into the North Sea via the River Wear (Fig. 1). The Wear has large stocks of Atlantic salmon and sea trout with annual resistivity (partial) counts of upstream adult migrants at Durham, 29 km from the sea, averaging 15,593 fish per year between 2007 and 2014 (Environment Agency, 2015). The Deerness drains a catchment of 52.4 km^2 , and is 16.3 km long, consisting largely of grassland to the west, and transitioning to woodland and lower-lying arable areas in the east. The brown trout population consists of multiple cohorts, with a mixture of resident and migratory adults, including anadromous trout (E. Winter, J. Tummers unpublished data).

The River Villestrup ($56^\circ 46' \text{ N}$, $9^\circ 55' \text{ E}$) is the primary freshwater source for the strongly brackish Mariager Fjord, ultimately exchanging with the Kattegat Sea on the east coast of Jutland, Denmark (Fig. 1). The river has a mean annual discharge of $1.1 \text{ m}^3 \text{ s}^{-1}$. The Villestrup is approximately

20 km long, has a typical width in the studied reach of 4-10 m and drains a catchment of 126 km². The average density of wild 0+trout is estimated at 125 per 100 m² (HELCOM, 2011). No stocking occurs. The inner fjord has salinities of 12-17 PSU in the upper 10 m of the water column used by trout, while deeper areas are more saline but often hypoxic. The shallow outer fjord has salinities of 20-25 PSU. The Villestrup joins near the junction between the inner and outer fjord areas, on the north shore (Fig. 1).

PIT tagging, recapture and telemetry

Trout in the Deerness system were captured for PIT-tagging using electric-fishing equipment at six sites dispersed over *ca.* 15 km of stream length (Fig. 1). Tagging occurred between 9 July and 12 September 2014 ($n = 643$), with a small number ($n = 23$) also tagged on 6 November 2014 (mean FL \pm SD of all trout = 151 mm \pm 23). In the Villestrup, trout were tagged on 26 September 2014 ($n = 490$; mean FL \pm SD = 147 mm \pm 27), using electric-fishing in a single reach *ca.* 8.5 km upstream of the river outlet (Fig. 1). Parr ≥ 120 mm (Larsen et al. 2013) and ≤ 250 mm from each study area were anaesthetised (Deerness: buffered MS-222, 100 mg L⁻¹; Villestrup: Benzocaine, 25 mg L⁻¹), weighed (to 0.1 g), measured (fork length, FL to 1 mm) and surgically implanted with a PIT-tag (Texas Instruments; model RI-TRP-RRHP, HDX, 134.2 kHz, length 23.1 mm, diameter 3.85 mm, weight 0.6 g in air). Tags and instruments were disinfected with 90% ethanol and air dried before use. Procedures were carried out by an experienced fish surgeon and following local animal welfare regulations. Following recovery, all individuals were returned to their site of capture. Recapture methodology used to investigate summer dispersal on the Deerness is presented in Online Resource 1.

Three pairs of stream-width swim-through half-duplex (HDX) PIT antennae were installed on a 5 km stretch of the lower Deerness (Fig. 1; Bolland et al. 2009), operational from 24 September 2014 at stations M2 and M3 and 13 October at station M1 (Fig. 1), until 31 May 2015. Due to occasional battery failure and a damaging high flow event in November causing severe loss of efficiency, the stations M1, M2 and M3 were operational 95.6, 98.1 and 98.1 % of the time, respectively. On the

Villestrup, a single pair of mains-powered antennae were placed 300 m upstream of the river outlet (Fig. 1), functional 88.7 % of the time from 26 September 2014 to 31 May 2015. Tags were detected by time-synchronised Master and Slave HDX readers (Texas Instruments SX2000; in-house build), interrogating the pairs of antennae eight times per second (Castro-Santos et al. 1996). Detection ranges between 20 and 80 cm were achieved and correct function on the Deerness was confirmed by passing a test tag through each antenna before and after each battery change (every 4 ± 2 days), and by more detailed range testing periodically. Each Villestrup antenna had a timed auto-emitter check tag (Oregon RFID). Additionally on the Villestrup, a Wolf-type trap (Wolf, 1951) situated directly downstream of the PIT antennae captured migrating smolts from 18 March 2015 until the end of the study. For the purpose of this study, and by reference to standard terminology applied elsewhere (e.g. Ibbotson et al. 2013), all fish detected prior to 1 February were labelled autumn migrants, while those detected from 1 February to 31 May were labelled spring smolts.

Theoretical antenna efficiencies of 99, 100 and 100% were achieved for stations M1-3, respectively, by routinely passing a test tag through the system. In practice, several factors influence a tag's probability of detection, including environmental conditions, tag velocity, tag orientation and the presence of other tags (Zydlewski et al. 2006; Burnett et al. 2013). A practical estimate of efficiency is, hence, given by the ratio of fish detected at a site that are known to have passed through (Zydlewski et al. 2006), and was estimated at 98.4% for M2. The efficiencies of M1 and M3 could not be estimated using this method, due to the absence of detection equipment downstream and the time lag between tagging events and the onset of monitoring, meaning the location of individuals prior to detection was uncertain. Using Zydlewski et al.'s (2006) method, the efficiency of the Villestrup station was estimated as 86.5% in spring, by identifying individuals caught in the downstream trap that were not detected by the PIT antennae.

Population density estimations and environmental monitoring

Quantitative estimates of trout densities ($n \cdot 100 \text{ m}^{-2}$) were made at each of the Deerness tagging sites, using a multiple-pass depletion method (Carle & Strub, 1978) in July 2014 (average triple-pass catchability 97.2% for \geq Age 1) and subsequently in March 2015 (average triple-pass catchability 98.4% for \geq Age 1). Developmental state of recaptured individuals (parr, smolt [including part-smolt] or adult) was also recorded, based on phenotypic characteristics (e.g. parr marks, body colour, body shape; Tanguy et al. 1994), in order to predict the seaward movement of certain individuals.

One logger (HOBO®; model U20-001-01; Onset Computer Corporation), situated at M3 on the Deerness, recorded temperature (accuracy $\pm 0.4 \text{ }^{\circ}\text{C}$) and water pressure (accuracy $\pm 0.6 \text{ kPa}$, converted to river level) at 15-min intervals throughout the study. On the Villestrup, temperature was measured at the upper antenna (Tinytag plus 2; model TGP-4017, www.geminidataloggers.com) and water level records were obtained from a fixed gauging station ca. 1 km upstream of the river outlet (Fig. 1).

Statistical analyses

The influence of environmental factors on the timing of downstream movement was analysed using generalised linear models (GLMs), comparing daily detection frequency with fluctuations in average daily water temperature ($^{\circ}\text{C}$), photoperiod, average daily water level (m) and the change in average daily water level (m). Initial Poisson GLMs revealed non-linear residual patterns and overdispersion, which was corrected for using the negative binomial distribution with a log-link function (Richards, 2008). All combinations of explanatory variables were trialled, but never utilising collinear factors (e.g. temperature and photoperiod). Following this, we explored the effects of subject-specific variables on the probability of tagged trout being detected as autumn or spring migrants. Binomial GLMs with a cloglog link function, due to asymmetry in the numbers of migrants and non-migrants, were constructed using combinations of the independent factors fish length (mm), mass (g) and Fulton's condition factor, but never with collinear variables (e.g. length and

mass). The additional variables tag-site density ($n \cdot 100m^{-2}$, summer 2014), tag-site distance upstream (km), and date of tagging were also analysed for subjects on the Deerness. Twelve Deerness fish, one a spring migrant, one an upstream migrant and ten undetected, and two Villestrup fish, one a spring migrant and one undetected, were omitted from this analysis due to a lack of mass, and hence condition, records. All analyses were conducted in R 3.1.1 (R Core Team, 2014) with use of the MASS package (Venables & Ripley, 2002). Model selection followed the minimisation of Akaike's information criterion (AIC) values, which represents the best compromise between lack of precision (too many parameters) and bias (too few parameters). Models within $\Delta 6$ AIC were retained, provided they were not increasingly complex versions of more efficient nested counterparts (outlined by Richards, 2008).

For further examination, we used independent-sample *t*-tests to compare the average length, mass, condition factor and net ground speed ($km \ day^{-1}$) of autumn and spring migrating individuals. Chi-squared tests with Yates' continuity correction were used to determine if the proportion of Deerness fish caught, inspected and categorised as smolts in March and subsequently detected downstream differed by tag site, and if the proportion of Deerness downstream migrants reaching M1 differed between autumn and spring.

Results

Site fidelity was strong during summer 2014, with 83% of recaptured Deerness trout parr ($n = 330$) travelling no further than 60 m and $< 0.01\%$ travelling over 200 m, from a previous known site of release (Online Resource 2). From autumn onwards, 140 (21.0%) of the 666 Deerness trout were PIT detected downstream of the site at which they were tagged and released, comprising 83 autumn migrants, 52 spring migrants and five individuals with activity spanning both periods (from here on labelled autumn migrants). Trout detected at M1 were assumed to have left the Deerness system, comprising 89 individuals in total (13.4%), 41 in autumn and 48 in spring. Migratory behaviour was observed throughout the period of study, with peaks of activity in October, November, March and

May (Fig. 2). Eleven upstream migrants from T1 to M3 were recorded, all occurring in autumn and none of which were detected leaving the stream. In March 2015, parr densities had decreased at all but one of the tagging sites (Table 1), by an overall average of 49%; a total of 50 tagged trout were recaptured in March across all release sites, of which 17 subsequently migrated downstream.

In the Villestrup, 195 of 490 tagged trout (39.8 %) were detected at the monitoring site and/or caught in the trap, comprising 49 autumn migrants, 136 spring migrants, and ten individuals with activity spanning both periods (from here on labelled autumn migrants). The Villestrup trout did not display a distinct autumnal peak of activity, however low levels of migratory behaviour were sustained throughout autumn and winter months. Activity of spring smolts peaked in late March, with continued high levels of movement throughout April and early May (Fig. 2).

Environmental regulators of downstream migratory behaviour

Mean daily water temperature and water level were retained in the best model for predicting autumn migrant activity in the Deerness (Table 2), both sharing significant positive relationships with daily detection frequency. No other combinations of variables were retained under the selection criteria, reflecting their poor explanatory power. In contrast, the change in mean daily water level was the best predictor of spring migrant activity in the Deerness, with a significant positive relationship. Two further models were retained utilising the variables water level and photoperiod, however greater ΔAIC values signified weaker explanatory power (Table 2). The best model predicting autumn migrant activity in the Villestrup retained both water level and temperature, but, in contrast to the Deerness, with a significant negative effect of temperature (Table 2). Under the selection criteria, five models were retained for predicting spring migrant activity in the Villestrup, with the best predictors being change in mean daily water level and water temperature in positive trends (Table 2). The removal of temperature produced a model with a ΔAIC of 0.3, indicating the explanatory power of temperature is low and water level on its own is a powerful predictor of daily smolt counts.

Phenotypic determinants of downstream migratory tendency

The best model describing autumn migrant probability in the Deerness retained fish mass and tag site distance upstream as predictive parameters, both with significant negative trends (Table 3; Fig. 3). Three alternative models utilising combinations of length, condition, distance and date can be found in Table 3. The best model for predicting spring migration retained subject mass, condition factor and tag site distance upstream. Mass and tag site distance shared significant negative relationships with probability of spring detection, while condition factor shared a positive relationship (Table 3; Fig. 3). Notably, tag site distance was retained in all models for the Deerness, suggesting it is a strong predictor of migratory tendency for all downstream migrating juveniles, but particularly for spring smolts, given the higher coefficient estimates. This is supported by significantly more smolts (captured, inspected and classified in March 2015) than expected being detected originating from T1, the most downstream tag site, relative to sites further upstream (Chi-square: $\chi^2_1 = 6.18$, $p = 0.013$). Autumn and spring migratory tendency in the Villestrup were best predicted by fish mass and condition, both sharing negative relationships in each scenario (Table 3; Fig. 3). For the autumn migrants, mass was a particularly strong predictor on its own, given the removal of condition to produce a model with a ΔAIC of only 0.7.

Comparisons of migratory phenotypes

There was no difference in length ($t_{138} = 0.22$, $p = 0.82$) or mass ($t_{137} = 0.39$, $p = 0.70$) at the time of tagging between autumn and spring migrants in the Deerness; however there was some evidence to suggest condition factor was greater in spring migrants ($t_{137} = 2.00$, $p = 0.048$; Fig. 3). Mean length and mass of upstream migrants was also significantly greater than that of all downstream migrants in the Deerness (Length: $t_{149} = 3.55$, $p < 0.001$; Mass: $t_{147} = 4.62$, $p < 0.001$; Fig. 3), although we found no difference in condition factor ($t_{147} = 0.05$, $p = 0.96$). There was no difference in length ($t_{193} = 0.12$, $p = 0.90$), mass ($t_{192} = 0.26$, $p = 0.79$) or condition factor ($t_{192} = 0.32$, $p = 0.75$; Fig. 3) between autumn and spring migrants in the Villestrup. Individual net ground speed of downstream migrants in the

Deerness varied dramatically from less than 1 to 88 km day⁻¹ throughout the study period, but on average, spring migrants travelled significantly faster (24.3 km day⁻¹) than autumn migrants (11.0 km day⁻¹) ($t_{127} = 3.82$, $p < 0.001$; Fig. 4). Additionally, downstream migrants were classified as stream 'emigrants' following a final detection at the most downstream monitoring site, ca. 700 m from the Deerness' confluence with the river Browney. A significantly greater proportion of spring migrants (84.6%) than autumn migrants (49.4%) became stream emigrants during the period in which they were first detected (Chi-square: $\chi^2_1 = 15.5$, $p < 0.001$). Five autumn non-emigrants were subsequently detected in the spring, four of which were then detected moving past M1.

Discussion

Extensive autumn downstream migrations were evident in juvenile brown trout (*Salmo trutta*) for both the Villestrup (Jutland, Denmark) and the Deerness (north-east England). Over 1.5 times the number of spring smolts were observed active between October and January in the Deerness. Moreover those leaving the system during autumn-winter represented 46% of all stream-emigrant juveniles in the study period. This contrasts the strong site fidelity observed during summer months and provides quantitative evidence of the protracted overwinter nature of the downstream migratory behaviour, broadly similar to that of the Marine Institute (2014), which states numbers of autumn migrating juveniles in the Burrishoole catchment, Ireland, have fluctuated between 18% and 57% of the total annual juvenile downstream migrant count, since 1982. Both the Burrishoole and Deerness catchments are characterised by mild, oceanic climates and frequent flow elevations following rainfall, possibly accounting for the prolonged 'autumn' downstream migration. Nevertheless, Deerness migrants cannot fully account for the marked reduction in parr densities at sites T2-T5 by March 2015, which must also reflect either local movement outside tagging sites, low overwinter survival, or a combination of both.

Probability of migration was higher in the Villestrup, but incidence of autumn migration was lower (25% of the total juvenile stream-emigrant count), yet remains higher than that reported by

Jonsson and Jonsson (2009) in Norway. Migrants in the Villestrup are assumed to enter the brackish Mariager Fjord soon after passage through the PIT monitoring station, however autumn migrants may have low gill Na^+K^+ -ATPase activity, as observed for autumn-emigrating *S. salar* juveniles (Riley et al. 2008). This could cause osmoregulatory stress, although salmonid parr have been known to reside in estuarine environments (Cunjak et al. 1989; Pinder et al. 2007). Conversely, individuals in the Deerness have much further to travel before reaching a saline environment, (meaning their migration strategy, i.e. anadromy or potamodromy, is unknown), and may explain why incidence of autumn migration is particularly high there. The recognition of an alternative juvenile downstream migratory phenotype is growing internationally and, hence, the potential contribution of these individuals to adult recruitment must be acknowledged.

The environmental factors regulating the phenology of downstream movement of salmonid smolts have been well studied, particularly the behavioural responses to water temperature, flow and light (e.g. Jonsson, 1991; Aarestrup et al. 2002; Aldvén et al. 2015). Stimuli for migration differ in their importance geographically between river systems and temporally between years (Hembre et al. 2001). This study suggests that water level (and hence river discharge) had the greatest influence on autumn migrant movement in both the Deerness and Villestrup, in accordance with Youngson et al. (1983), Jonsson and Jonsson (2002) and Holmes et al. (2014). This is not surprising, given the opportunity to minimise the energetic costs of migration, while high turbidity may offer greater protection from predators (Hvidsten & Hansen, 1989). The change in average daily water level was a better predictor of movement of spring migrants in both the Deerness and Villestrup. This suggests smolts in both systems are particularly receptive to dynamic fluctuations in the hydrograph, similar to Carlsen et al.'s (2004) conclusions that migrating juveniles can anticipate floods. Importantly, all models retained in Table 2 utilised an environmental variable associated with river discharge, suggesting it is a central migratory stimulus for both populations.

Autumn migrants in the Deerness and smolts in the Villestrup responded positively to higher temperature, analogous to results obtained by Jonsson and Ruud-Hansen (1985). Smolt activity in the Deerness was better predicted by photoperiod, which is known to regulate physiological changes associated with the parr-smolt transformation (Björnsson et al. 2011). The probability of autumn migration in the Villestrup shared a significant negative relationship with temperature (see also Jonsson & Jonsson, 2002), yet previous studies reporting increased migrant activity at cold temperatures are usually associated with ice melt (Hesthagen & Garnås, 1986; Carlsen et al. 2004). One possibility for the observed pattern is that peak flows may have coincided with low temperatures in winter, demonstrating a degree of behavioural independence with regard to seasonal variables. Also, the Villestrup is spring-fed, meaning temperatures are comparatively more stable to those of the Deerness. The relative importance of temperature, photoperiod and river discharge as migratory triggers may, however, fluctuate between years (Jensen et al. 2012; Aldvén et al. 2015), depending on precipitation and rate of temperature change. Longer-term and experimental studies in either system could reveal variability and mechanisms in the effect of environmental cues.

The probability of an individual conducting autumn or spring migration past fixed points on the Deerness decreased significantly in an upstream direction. Ibbotson et al. (2013) proposed a re-distribution of autumn migrating *S. salar* parr in a downstream direction was responsible for similar findings, rather than a targeted migration. One explanation for this may be to consider the potential disturbance caused by environmental events such as high autumn-winter flows. Territorial behaviour may be disrupted, initiating a re-establishment of dominance hierarchies and promoting the downstream displacement of subordinate individuals. While this may hold true for autumn-winter migrants, it is unlikely to be the case for spring smolts, but for which we found a reduced probability of detection from tag sites further upstream. Moreover, there appears no trend in the percentage of recaptured residents with distance upstream (Table 1), which would be expected if migration tendency decreased in an upstream direction. We, therefore, suggest a cumulative increase in

mortality probability with increasing distance upstream for both autumn and spring migrants, for example due to anthropogenic obstruction during migration, greater energetic costs or exposure to predators (Aarestrup & Koed, 2003; Gauld et al. 2013).

In early studies it was hypothesised that anadromous salmonid juveniles migrate at the first opportunity after reaching a threshold size (Elson, 1957; Fahy, 1985), yet for brown trout, smolt length can vary from <100 mm to >200 mm within the same river (Økland et al. 1993). Fast-growing individuals smolt at a younger age and smaller size than their slow-growing counterparts (Forseth et al. 1999). In the Deerness and Villestrup, autumn and spring migratory tendency were negatively affected by body mass, which was always a better predictor than body length. This may be counterintuitive, considering migration and post-migration survival is thought to be positively size-dependent (Bohlin et al. 1993), but could suggest migrating juveniles in these river systems were energetically constrained. Migration has been described as a biological response to adversity (Taylor & Taylor, 1977), and in the Deerness (autumn migrants) and Villestrup (all migrants) migratory tendency was negatively correlated with fish condition at tagging, which could result from adversity in the form of poor growth opportunities. Notably, the optimal size at migration in the Deerness and Villestrup systems could be lower than the size range of individuals sampled such that some of the tagged trout, especially males, may have been maturing.

The size of autumn and spring migrants did not differ at the time of tagging in either the Deerness or Villestrup, as found by Ibbotson et al. (2013) for *S. salar*. In contrast, some found autumn migrants to be significantly larger than spring migrants or residents of the same year-class, possibly indicating constrained habitat availability for faster-growing individuals or achievement of a high energy store status (Huntingford et al. 1992; Holmes et al. 2014). This hypothesis is not supported by our results. In the Deerness, spring migrants were of better condition than autumn migrants at the time of tagging and autumn and spring migratory tendencies shared opposed relationships with individual body condition. This may have been a result of competition, such that

subordinate individuals with lower body condition were displaced by their dominant counterparts, either preferentially leaving a low-growth potential environment or forced out of refugia with an overwinter reduction in stream carrying capacity (see Keeley, 2001). Migration in brown trout can be regulated by food availability (Wysujack et al. 2008), therefore autumn migrants could result from a competition-induced lack of resources for subordinates. On the contrary, we found no evidence to suggest the density of trout Age 1+ and older at each Deerness site could predict the proportion of autumn migrants. However, caution is needed, since local habitat has been shown to influence the autumnal movement of Atlantic salmon parr, perhaps due to differences in the proximity of overwintering habitat (Ibbotson et al. 2013).

Autumn upstream migrants in the Deerness, characteristic of precocious parr maturation (McCormick et al. 1998), were larger than downstream migrants. Precocious parr are predominantly males and the anadromous emigrants' sex ratio is typically heavily skewed towards females (Klemetsen et al. 2003) although autumn downstream migration has been linked to the reproductive motivations of precocious parr (Buck and Youngson, 1982). Fish sex was unknown in this study, although 22.5% of tagged and untagged parr morphotypes ≥ 120 mm, sampled on the Deerness in November 2014 were spermiating males (E. Winter, unpublished data). Cheap molecular methods for the sexing of juvenile salmonids from tissue samples are now available (Quéméré et al. 2014) and will aid sex-specific interpretation of movement patterns in parr morphotype salmonids.

Questions regarding the ecological significance of autumn migration of juvenile salmonids remain. Increased plasma thyroxine levels in autumn migrants are suggestive of a physiologically mediated migration (Riley et al. 2008; Zydlewski et al. 2005), however the behavioural motivations of autumn and spring migrants may differ, since autumn-migrating parr are not physiologically adapted for seawater entry (Riley et al. 2008). Deerness spring migrants travelled at a greater net ground speed, at rates similar to those obtained by Aarestrup et al. (2002) for radio tagged trout smolts, and were more likely to become stream emigrants than autumn migrants. While antennae

malfunction during a major spate in November 2014 may be partially responsible for a lack of autumn detections at M1, these results suggest the movements of Deerness autumn individuals are not exclusively marine-targeted. In the Villestrup it seems likely that downstream-migrating parr recorded at the PIT station at the bottom of the river subsequently enter the Mariager Fjord, though they could remain in the river outlet. It would, therefore, be valuable to determine the range of salinities and temperatures to which juvenile autumn emigrants are exposed and their survival in relation to their physiological readiness for seawater transition.

In conclusion, this study provides quantitative evidence for considerable autumn and winter downstream migration of juvenile brown trout in the rivers Deerness and Villestrup, along with correlative information on factors regulating behaviour, to parallel the wealth of knowledge concerning spring smolt migration. We propose different behavioural motivations for autumn and spring migrants, based on individual condition, the proportion of emigrants and rates of migration for each group in the Deerness. We emphasise the dynamic behavioural nature of the brown trout and suggest the autumn-migratory phenotype represents an important avenue within the migration continuum concept (Cucherousset et al. 2005; Dodson et al. 2013; Boel et al. 2014), dependent on environmental and physiological factors relating to individual fitness. An in-depth, experimental evaluation of the temporal, spatial and genetic variability of the extent of autumn migration of trout, and its influence on subsequent life history traits, would be beneficial to future conservation and management plans.

Acknowledgements

We are grateful to Sean Twiss and Amy Bishop for assistance with statistical analyses. Thanks also go to landowners for granting access to the Deerness sites and to Elana Hobkirk for assistance with fieldwork.

References

403 Aarestrup, K. & A. Koed, 2003. Survival of migrating sea trout (*salmo trutta*) and Atlantic salmon
 404 (*Salmo salar*) smolts negotiating weirs in small Danish rivers. *Ecology of Freshwater Fish* 12: 169-176.

405 Aarestrup, K., C. Nielsen & A. Koed, 2002. Net ground speed of downstream migrating radio-tagged
 406 Atlantic salmon (*Salmo salar* L.) and brown trout (*Salmo trutta* L.) smolts in relation to
 407 environmental factors. *Hydrobiologia* 483: 95-102.

408 Aarestrup, K., H. Baktoft, A. Koed, D. del Villar-Guerra & E. B. Thorstad, 2014. Comparison of the
 409 riverine and early marine migration behaviour and survival of wild and hatchery-reared sea trout
 410 *Salmo trutta* smolts. *Marine Ecology Progress Series* 496: 197-206.

411 Aldvén, D., E. Degerman & J. Höjesjö, 2015. Environmental cues and downstream migration of
 412 anadromous brown trout (*Salmo trutta*) and Atlantic salmon (*Salmo salar*) smolts. *Boreal*
 413 *Environment Research* 20: 35-44.

414 Björnsson, B. T., S. O. Stefansson & S. D. McCormick, 2011. Environmental endocrinology of salmon
 415 smoltification. *General and Comparative Endocrinology* 170: 290-298.

416 Boel, M., K. Aarestrup, H. Baktoft, T. Larsen, S. S. Madsen, H. Malte, C. Skov, J. C. Svendsen & A.
 417 Koed, 2014. The physiological basis of the migration continuum in brown trout (*Salmo trutta*).
 418 *Physiological and Biochemical Zoology* 87: 334-345.

419 Bohlin, T., C. Dellefors & U. Faremo, 1993. Optimal time and size for smolt migration in wild sea trout
 420 (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* 50: 224-232.

421 Bohlin, T., J. Pettersson & E. Degerman, 2001. Population density of migratory and resident brown
 422 trout (*Salmo trutta*) in relation to altitude: evidence for a migration cost. *Journal of Animal Ecology*
 423 70: 112-121.

424 Bolland, J. D., I. G. Cowx & M. C. Lucas, 2009. Dispersal and survival of stocked cyprinids in a small
 425 English river: comparison with wild fish using a multi-method approach. *Journal of Fish Biology* 74:
 426 2313-2328.

427 Buck, R. J. G. & A. F. Youngson, 1982. The downstream migration of precociously mature Atlantic
 428 salmon, *Salmo salar* L. parr in autumn; its relation to the spawning migration of mature adult fish.
 429 *Journal of Fish Biology* 20: 279-288.

430 Burnett, N. J., K. M. Stamplecoskie, J. D. Thiem & S. J. Cooke, 2013. Comparison of detection
 431 efficiency among three sizes of half-duplex passive integrated transponders using manual tracking
 432 and fixed antenna arrays. *North American Journal of Fisheries Management* 33: 7-13.

433 Carle, F. L., & M. R. Strub, 1978. A new method for estimating population size from removal data.
 434 *Biometrics* 34: 621-630.

435 Carlsen, K. T., O. K. Beg, B. Finstad & T. G. Heggberget, 2004. Diel periodicity and environmental
 436 influence on the smolt migration of Arctic char, *Salvelinus alpinus*, Atlantic salmon, *Salmo salar*, and
 437 sea trout, *Salmo trutta*, in northern Norway. *Environmental Biology of Fishes* 70: 403-413.

438 Castro-Santos, T., A. Haro & S. Walk, 1996. A passive integrated transponder (PIT) tag system for
 439 monitoring fishways. *Fisheries Research* 28: 253-261.

440 Chapman, B. B., C. Bronmark, J. A. Nilsson & L. A. Hansson, 2011. The ecology and evolution of
 441 partial migration. *Oikos* 120: 1764-1775.

442 Cucherousset, J., D. Ombredane, K. Charles, F. Marchand & J.-L. Baglinière, 2005. A continuum of life
 443 history tactics in a brown trout (*Salmo trutta*) population. *Canadian Journal of Fisheries and Aquatic*
 444 *Sciences* 62: 1600-1610.

445 Cunjak, R. A., 1992. Comparative feeding, growth and movement of Atlantic salmon (*Salmo salar*)
 446 parr from riverine and estuarine environments. Ecology of Freshwater Fish 1: 26-34.

447 Cunjak, R. A., E. M. P. Chadwick & M. Shears, 1989. Downstream movements and estuarine
 448 residence by Atlantic salmon parr (*Salmo salar*). Canadian Journal of Fisheries and Aquatic Sciences
 449 46: 1466-1471.

450 del Villar-Guerra, D., K. Aarestrup, C. Skov & A. Koed, 2014. Marine migrations in anadromous brown
 451 trout (*Salmo trutta*). Fjord residency as a possible alternative in the continuum of migration to the
 452 open sea. Ecology of Freshwater Fish 23: 594-603.

453 Dodson, J. J., N. Aubin-Horth, V. Thériault & D. J. Páez, 2013. The evolutionary ecology of alternative
 454 migratory tactics in salmonid fishes. Biological Reviews 88: 602-625.

455 Elliott, J. M., 1986. Spatial distribution and behavioural movements of migratory trout *Salmo trutta*
 456 in a Lake District stream. Journal of Animal Ecology 56: 83-98.

457 Elliott, J. M., 1994. Quantitative ecology and the brown trout. Oxford University Press, Oxford.

458 Elson, P. F., 1957. The importance of size in the change from parr to smolt in Atlantic salmon.
 459 Canadian Fish Culturist 21: 1-6.

460 Environment Agency, 2015. River Wear monthly upstream fish counts.
 461 <https://www.gov.uk/government/statistics/river-wear-monthly-upstream-fish-counts>

462 Fahy, E., 1985. Cyclic fluctuations in the abundance of trout (*Salmo trutta* L.). Archiv fuer
 463 Hydrobiologie Supplement 70: 404-428.

464 Forseth T., T. F. Næsje, B. Jonsson & K. Hårsaker, 1999. Juvenile migration in brown trout: a
 465 consequence of energetic state. Journal of Animal Ecology 68: 783-793.

466 Gauld, N. R., R. Campbell & M. C. Lucas, 2013. Reduced flows impact salmonid smolt emigration in a
 467 river with low-head weirs. *Science of the Total Environment* 458-460: 435-443.

468 HELCOM, 2011. Sea Trout and Salmon Populations and Rivers in Denmark – HELCOM assessment of
 469 salmon (*Salmo salar*) and sea trout (*Salmo trutta*) populations and habitats in rivers flowing to the
 470 Baltic Sea. Baltic Sea Environment Proceedings No. 126B.

471 Hembre, B., J. V. Arnekleiv & J. H. L'Abée-Lund, 2001. Effects of water discharge and temperature on
 472 the seaward migration of anadromous brown trout, *Salmo trutta*, smolts. *Ecology of Freshwater Fish*
 473 10: 61-64.

474 Hesthagen, T. & E. Garnås, 1986. Migration of Atlantic salmon smolts in River Orkla of central
 475 Norway in relation to management of a hydroelectric station. *North American Journal of Fisheries*
 476 *Management* 6: 376-382.

477 Holmes, R., J. W. Hayes, W. Jiang, A. Quarterman & L. N. Davey, 2014. Emigration and mortality of
 478 juvenile brown trout in a New Zealand headwater tributary. *Ecology of Freshwater Fish* 23: 631-643.

479 Huntingford, F. A., J. E. Thorpe, C. G. de Leaniz & D. W. Hay, 1992. Patterns of growth and smolting in
 480 autumn migrants from a Scottish population of Atlantic salmon, *Salmo salar* L. *Journal of Fish Biology*
 481 41:43-51.

482 Hvidsten, N. A. & L. P. Hansen, 1989. Increased recapture rate of adult Atlantic salmon, *Salmo salar*
 483 L., stocked as smolts at high water discharge. *Journal of Fish Biology* 32: 153-154.

484 Ibbotson, A. T., W. D. Riley, W. R. C. Beaumont, A. C. Cook, M. J. Ives, A. C. Pinder & L. J. Scott, 2013.
 485 The source of autumn and spring downstream migrating juvenile Atlantic salmon in a small lowland
 486 river. *Ecology of Freshwater Fish* 22: 73-81.

487 Jensen, A. J., B. Finstad, P. Fiske, N. A. Hvidsten, A. H. Rikardsen & L. Saksgård, 2012. Timing of smolt
 488 migration in sympatric populations of Atlantic salmon (*Salmo salar*), brown trout (*Salmo trutta*), and
 489 Arctic char (*Salvelinus alpinus*). Canadian Journal of Fisheries and Aquatic Sciences 69: 711-723.

490 Jonsson, B., 1985. Life history patterns of freshwater resident and sea-run migrant brown trout in
 491 Norway. Transactions of the American Fisheries Society 114: 182-194.

492 Jonsson, B. & N. Jonsson, 2009. Migratory timing, marine survival and growth of anadromous brown
 493 trout *Salmo trutta* in the River Imsa, Norway. Journal of Fish Biology 74: 621-638.

494 Jonsson, B. & N. Jonsson, 2011. Ecology of Atlantic salmon and Brown Trout: Habitat as a template
 495 for life histories. Fish and Fisheries Series 33, Springer, Dordrecht, the Netherlands.

496 Jonsson, B., & J. Ruud-Hansen, 1985. Water temperature as the primary influence on timing of
 497 seaward migrations of Atlantic salmon (*Salmo salar*) smolts. Canadian Journal of Fisheries and
 498 Aquatic Sciences 42: 593-595.

499 Jonsson, N., 1991. Influence of water flow, water temperature and light on fish migration in rivers.
 500 Nordic Journal of Freshwater Research 66: 20-35.

501 Jonsson, N. & B. Jonsson, 2002. Migration of anadromous brown trout in a Norwegian river.
 502 Freshwater Biology 47: 1391-1401.

503 Jonsson, N. & B. Jonsson, 2014. Time and size at seaward migration influence the sea survival of
 504 Atlantic salmon (*Salmo salar* L.). Journal of Fish Biology 84: 1457-1473.

505 Keeley, E. R., 2001. Demographic responses to food and space competition by juvenile steelhead
 506 trout. Ecology 82: 1247-1259.

507 Klemetsen, A., P. A. Amundsen, J. B. Dempson, B. Jonsson, N. Jonsson, M. F. O'Connell & E.
 508 Mortensen, 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr
 509 *Salvelinus alpinus* (L.): a review of aspects of their life histories. Ecology of Freshwater Fish 12: 1-59.
 510 Larsen, M. H., A. N. Thorn, C. Skov & K. Aarestrup, 2013. Effects of passive integrated transponder
 511 tags on survival and growth of juvenile Atlantic salmon *Salmo salar*. Animal Biotelemetry 1: 1-7.
 512 Lucas, M. C. & E. Baras, 2001. Migration of Freshwater Fishes. Blackwell Science Ltd, Oxford.
 513 Marine Institute 2014. Newport Research Facility, Annual Report No. 58, 2013.
 514 McCormick, S. D., L. P. Hansen, T. P. Quinn & R. L. Saunders, 1998. Movement, migration, and
 515 smolting of Atlantic salmon (*Salmo salar*). Canadian Journal of Fisheries and Aquatic Sciences 55: 77-
 516 92.
 517 McGinnity, P., E. de Eyto, T. F. Cross, J. Coughlan, K. Whelan & A. Ferguson, 2007. Population specific
 518 smolt development, migration and maturity schedules in Atlantic salmon in a natural river
 519 environment. Aquaculture 273: 257-268.
 520 Nathan, R., W. M. Getz, E. Revilla, M. Holyoak, R. Kadmon, D. Saltz & P. E. Smouse, 2008. A
 521 movement ecology paradigm for unifying organismal movement research. PNAS 105: 19052-19059.
 522 Økland, F., B. Jonsson, A. J. Jensen & L. P. Hansen, 1993. Is there a threshold size regulating seaward
 523 migration of brown trout and Atlantic salmon? Journal of Fish Biology 42: 541-550.
 524 Pinder, A. C., W. D. Riley, A. T. Ibbotson & W. R. C. Beaumont, 2007. Evidence for an autumn
 525 downstream migration and the subsequent estuarine residence of 0+ year juvenile Atlantic salmon,
 526 *Salmo salar* L., in England. Journal of Fish Biology 71: 260-264.

527 Quéméré, E., C. Perrier, A.-L. Besnard, G. Evanno, J.-L. Baglinière, Y. Guiguen & S. Launey, 2014. An
 528 improved PCR-based method for faster sex determination in brown trout (*Salmo trutta*) and Atlantic
 529 salmon (*Salmo salar*). Conservation Genetics Resources, 6: 825-827.

530 R Core Team, 2014. R: A language and environment for statistical computing. R Foundation for
 531 Statistical Computing, Vienna, Austria, <http://www.R-project.org/>.

532 Richards, S. A., 2008. Dealing with overdispersed count data in applied ecology. Journal of Applied
 533 Ecology, 45: 218–227.

534 Riddell, B. E., & W. C. Leggett, 1981. Evidence of an adaptive basis for geographic variation in body
 535 morphology and time of downstream migration of juvenile Atlantic salmon (*Salmo salar*). Canadian
 536 Journal of Fisheries and Aquatic Sciences 38: 308-320.

537 Riley, W. D., A. T. Ibbotson, N. Lower, A. C. Cook, A. Moore, S. Mizuno, A. C. Pinder, W. R. C.
 538 Beaumont & L. Privitera, 2008. Physiological seawater adaptation in juvenile Atlantic salmon (*Salmo*
 539 *salar*) autumn migrants. Freshwater Biology 53: 745-755.

540 Riley, W. D., A. T. Ibbotson & W. R. C. Beaumont, 2009. Adult returns from Atlantic salmon, *Salmo*
 541 *salar*, parr autumn migrants. Fisheries Management and Ecology 16: 75-76.

542 Taal, I., M. Kesler, L. Saks, M. Rohtla, A. Verliin, R. Svirsden, K. Jurgens, M. Vetemaa & T. Saat, 2014.
 543 Evidence for an autumn downstream migration of Atlantic salmon *Salmo salar* (Linnaeus) and brown
 544 trout *Salmo trutta* (Linnaeus) parr to the Baltic Sea. Helgoland Marine Research 68: 373-377.

545 Tanguy, J., D. Ombredane, J. Bagliniere & P. Prunet, 1994. Aspects of parr-smolt transformation in
 546 anadromous and resident forms of brown trout (*Salmo trutta*) in comparison with Atlantic salmon
 547 (*Salmo salar*). Aquaculture 121: 51-63.

548 Taylor, L. R. & R. A. J. Taylor, 1977. Aggregation, migration and population mechanics. *Nature* 265:
 549 415-421.

550 Venables, W. N., & B. D. Ripley, 2002. *Modern Applied Statistics with S*. Fourth Edition. Springer,
 551 New York. ISBN 0-387-95457-0

552 Wolf, P. A., 1951. A trap for the capture of fish and other organisms moving downstream.
 553 *Transactions of the American Fisheries Society* 80: 41-45.

554 Wysujack, K., L. A. Greenberg, E. Bergman & I. C. Olsson, 2008. The role of the environment in partial
 555 migration: food availability affects the adoption of a migratory tactic in brown trout *Salmo trutta*.
 556 *Ecology of Freshwater Fish* 18: 52-59.

557 Youngson, A. F., R. J. G. Buck, T. H. Simpson & D. W. Hay, 1983. The autumn and spring emigrations
 558 of juvenile Atlantic salmon, *Salmo salar* L., from the Girnock Burn, Aberdeenshire, Scotland:
 559 environmental release of migration. *Journal of Fish Biology* 23: 625-639.

560 Zydlewski, G. B., A. Haro & S. D. McCormick, 2005. Evidence for cumulative temperature as an
 561 initiating and terminating factor in downstream migratory behaviour of Atlantic salmon (*Salmo*
 562 *salar*) smolts. *Canadian Journal of Fisheries and Aquatic Sciences* 62: 68-78.

563 Zydlewski, G. B., G. Horton, T. Dubreuil, B. Lecther, S. Casey & J. Zydlewski, 2006. Remote monitoring
 564 of fish in small streams: a unified approach using PIT tags. *Fisheries* 31: 492-502.

565 Figure legends

566 Fig. 1 Map of the Deerness and Villestrup study areas, showing the locations of tagging sites, fixed
567 PIT readers and environmental monitoring stations with stars, thick lines and open circles,
568 respectively. On the Deerness, an environmental monitoring station was also placed at M3. Inset
569 maps show the location of the study areas nationally.

570 Fig. 2 Daily detection frequency of PIT tagged trout parr detected downstream of the site at which
571 they were tagged and released on the Deerness (a) and Villestrup (c). Date of first detection only is
572 displayed. Shaded regions represent periods of minimal or no detection efficiency due to PIT
573 antennae malfunction. Individuals detected prior to 1 Feb were labelled autumn migrants and those
574 detected from 1 Feb labelled spring smolts following published convention (Ibbotson et al. 2013). In
575 addition, a downstream trap was operational from 18 March on the Villestrup, indicated by the
576 arrow on panel (c). Average daily water temperature (solid lines) and average daily water level
577 (dotted lines) are also displayed for the Deerness (b) and Villestrup (d)

578 Fig. 3 The mean length, mass and condition factor at tagging (\pm SEM) of undetected trout, autumn
579 downstream migrants, spring downstream migrants and upstream migrants in the Deerness (top
580 row) and the Villestrup (bottom row)

581 Fig. 4 The mean net ground speed (\pm SEM) of autumn and spring migrants in the Deerness.

582




Fig. 1




Fig. 2




Fig. 3




Fig. 4

598 Table 1: Details of tag site-specific variables on the Deerness, along with results of PIT telemetry and March recapture surveys by site.

Tag Site	Distance from river mouth (km)	Density '14 (n·100m ⁻²)	Density '15 (n·100m ⁻²)	%Δ Density	No. fish tagged	% Migrated Downstream	% Recaptured residents	% Loss from mortality or local dispersal
T1	4.2	12.0	14.3	+19.2	214	33.6	4.2	62.2
T2	6.2	9.2	3.7	-59.8	99	10.1	6.1	83.8
T3	8.6	12.2	1.9	-84.4	91	28.6	3.3	68.1
T4	10.8	9.5	1.5	-84.2	55	18.2	7.3	74.5
T5	11.8	12.2	2.3	-81.1	140	12.1	2.1	85.8
T6	14.5	7.7	7.5	-2.6	67	7.5	10.4	82.1

599

600

601 Table 2: Generalised linear model outputs of migration phenology analyses. All retained models within 6 Δ AIC are displayed. Significant variables are in
602 bold, with values for predictor variables representing coefficient estimates.

Model	AICc	Δ AICc	df	Intercept	Water Level	Δ Water Level	Water Temperature	Photoperiod
Deerness Autumn								
1	202.3	0.0	4	-12.00	33.30		0.24	
Deerness Spring								
1	166.8	0.0	3	-1.76		22.83		
2	168.3	1.5	4	-9.51	20.24			0.004
3	169.1	2.3	3	-5.84	17.35			
Villestrup Autumn								
1	207.7	0.0	4	-20.80	10.83		-0.34	
2	208.6	0.9	4	-18.23	11.02			-0.011
Villestrup Spring								
1	311.7	0.0	4	-1.87		27.04	0.20	
2	312.0	0.3	3	-0.26		21.68		
3	316.1	4.4	4	-29.64	13.77			0.004
4	316.4	4.7	4	-34.35	16.62		0.28	
5	317.3	5.6	3	-17.57	9.10			

603
604

605 Table 3: Generalised linear model outputs of migration tendency analyses. All retained models within 6 Δ AIC are displayed. Significant variables are in bold,
606 with values for predictor variables representing coefficient estimates.

Model	AICc	ΔAICc	df	Intercept	Length	Weight	Condition	Density	Distance	Days
Deerness Autumn										
1	487.4	0.0	3	0.26		-0.032			-0.136	0.006
2	489.7	2.3	5	4.29	-0.023		-1.86		-0.146	
3	489.7	2.3	4	3.93	-0.022		-1.42		-0.137	
4	491.3	3.9	3	2.48	-0.022				-0.143	
Deerness Spring										
1	340.7	0.0	4	-1.35		-0.041	1.79		-0.178	
2	341.0	0.3	3	3.42	-0.029				-0.178	
3	343.0	2.3	3	0.24		-0.035			-0.164	
Villemstrup Autumn										
1	353.9	0.0	3	2.15		-0.023	-3.33			
2	354.6	0.7	2	-1.21		-0.024				
3	356.0	2.1	3	4.34	-0.014		-4.19			
4	358.4	4.5	2	-0.002	-0.014					
5	359.3	5.4	2	2.34			-4.22			
Villemstrup Spring										
1	527.0	0.0	3	4.70		-0.026	-4.42			
2	532.6	5.6	3	7.40	-0.017		-5.47			

Online Resource 1: Investigating the phenology of downstream migration in juvenile trout.

Methodology of Deerness recapture surveys in summer 2014, used to investigate summer-time dispersal of trout parr.

Following the initial PIT-tagging of trout parr at sites along the Deerness, recapture sessions during the summer of 2014 enabled the re-sampling of individuals to determine their level of movement. Three single-pass electrofishing surveys were conducted at each site in contiguous stop-netted 20 m stream sections over a period of *ca.* two months (early July to mid September 2014). The distance surveyed was increased in each session from initially 100 m to finally 200 m upstream and downstream of each starting point. During the first recapture session at sites T2, T3 and T4, and during the first two recapture sessions at sites T1, T5 and T6, any untagged individuals captured were tagged and subsequently released back into the population. The frequencies of maximum longitudinal distances covered by each recaptured fish from a previous site of release showed a typical inverse-power distribution. Following the methods of Bubb et al. (2004), linear transformation using a double-In plot enabled regression analysis. Upstream and downstream dispersal were analysed separately and the regression lines compared. To maximise sample size, the recorded dispersal ranges were combined from all study sites.

References

Bubb, D. H., T. J. Thom & M. C. Lucas, 2004. Movement and dispersal of the invasive signal crayfish *Pacifastacus leniusculus* in upland rivers. *Freshwater Biology*, 49: 357-368.

Online Resource 2: Investigating the phenology of downstream migration in juvenile trout.

Results and analysis of the Deerness recapture surveys of trout parr dispersal in summer 2014.

A total number of 330 fish were recaptured during single pass recapture surveys on the Deerness between late July and mid-September 2014. The maximum distance covered by any one trout tagged and recaptured within the same Deerness study site, July to September 2014, was 260 m, however many recaptured individuals (50.2%) remained within the 20m zone in which they were first released, and the majority (83.0%) travelled no further than 60m from previous known locations (Fig. S1). The probability, M , of moving a dispersal distance of D m was described by an inverse-power function using the inverse cumulative proportion of dispersers in each 20 m sample zone from the site of release (Bubb et al. 2004):

$$M = CD^{-n}$$

C and n are scaling constants. A highly significant negative relationship between $\ln M$, both upstream and downstream, and $\ln D$ was found (Upstream: $F_{1,7} = 82.7$, $p < 0.001$, $R^2 = 0.92$; Downstream: $F_{1,10} = 121.7$, $p < 0.001$, $R^2 = 0.92$; Fig. S2) under the equations:

$$\ln M \text{ (upstream)} = 7.25 - 2.26(\ln D)$$

$$\ln M \text{ (downstream)} = 7.22 - 2.21(\ln D)$$

No significant difference between the gradients of the two regression lines was found ($t_{20} = -0.18$, $p = 0.86$), meaning trout were equally likely to disperse upstream and downstream (Fig. S2).




Fig. S1: Frequency distribution of the maximum longitudinal movements made by trout from a previous release site (top), with upstream and downstream movements separated (bottom), all study sites combined.




Fig. S2: Double-In plot of the inverse cumulative proportion of dispersers in upstream (grey) and downstream (blue) directions.

647

648 **References**

649 Bubb, D. H., T. J. Thom & M. C. Lucas, 2004. Movement and dispersal of the invasive signal crayfish
 650 *Pacifastacus leniusculus* in upland rivers. *Freshwater Biology*, 49: 357-368.

